

CENOZOIC GEOLOGY SOUTHWEST USA

THE CENOZOIC GEOLOGY OF THE SOUTHWEST USA REPRESENTS A DRAMATIC AND DYNAMIC CHAPTER IN EARTH'S HISTORY, PROFOUNDLY SHAPING THE ICONIC LANDSCAPES WE SEE TODAY. THIS ERA, SPANNING THE LAST 66 MILLION YEARS, WITNESSED IMMENSE GEOLOGICAL FORCES AT PLAY, FROM THE RIFTING AND VOLCANISM THAT CREATED VAST BASINS AND RANGES TO THE EROSIONAL SCULPTORS THAT CARVED CANYONS AND MESAS. UNDERSTANDING THE CENOZOIC GEOLOGICAL PROCESSES IN THIS REGION IS KEY TO APPRECIATING ITS RICH MINERAL RESOURCES, ITS UNIQUE ECOSYSTEMS, AND THE CHALLENGES OF LIVING IN AN ACTIVE GEOLOGICAL ENVIRONMENT. THIS ARTICLE WILL DELVE INTO THE INTRICATE TAPESTRY OF CENOZOIC EVENTS, EXPLORING ITS MAJOR EPOCHS, TECTONIC FORCES, VOLCANIC ACTIVITY, SEDIMENTARY DEPOSITION, AND THE RESULTING LANDFORMS THAT DEFINE THE SOUTHWEST.

TABLE OF CONTENTS

THE DAWN OF THE CENOZOIC: PALEOGENE AND NEOGENE FOUNDATIONS

TECTONIC DRIVERS OF CENOZOIC SOUTHWEST GEOLOGY

VOLCANIC PROVINCES AND THEIR ENDURING LEGACY

SEDIMENTARY BASINS AND DEPOSITIONAL ENVIRONMENTS

EROSIONAL SCULPTURES: THE ARID SOUTHWEST'S MASTERPIECES

SIGNIFICANT CENOZOIC GEOLOGICAL FEATURES OF THE SOUTHWEST

THE DAWN OF THE CENOZOIC: PALEOGENE AND NEOGENE FOUNDATIONS

THE CENOZOIC ERA COMMENCED WITH THE PALEOGENE PERIOD, A TIME OF SIGNIFICANT RECOVERY AND DIVERSIFICATION OF LIFE FOLLOWING THE CRETACEOUS-PALEOGENE EXTINCTION EVENT. IN THE SOUTHWEST USA, THIS PERIOD WAS CHARACTERIZED BY THE WANING INFLUENCE OF EARLIER TECTONIC REGIMES AND THE NASCENT STAGES OF THE LARAMIDE OROGENY'S FAR-REACHING EFFECTS. WHILE THE MAJOR MOUNTAIN-BUILDING ASSOCIATED WITH THE LARAMIDE PEAKED CLOSER TO THE BOUNDARY OF THE MESOZOIC, ITS COMPRESSIONAL FORCES CONTINUED TO RIPPLE THROUGH THE REGION, INFLUENCING CRUSTAL DEFORMATION AND THE INITIAL DEVELOPMENT OF SEDIMENTARY BASINS. SEDIMENTARY ROCKS FROM THE PALEOGENE IN THE SOUTHWEST OFTEN REFLECT EXTENSIVE FLUVIAL AND LACUSTRINE (LAKE) ENVIRONMENTS, DEPOSITING SANDS, SILTS, AND CLAYS THAT WOULD LATER BE LITHIFIED INTO FORMATIONS LIKE THE GREEN RIVER FORMATION, FAMOUS FOR ITS FOSSILS AND OIL SHALE.

FOLLOWING THE PALEOGENE, THE NEOGENE PERIOD USHERED IN FURTHER DRAMATIC TRANSFORMATIONS. THIS ERA SAW THE INTENSIFICATION OF BASIN AND RANGE EXTENSION ACROSS MUCH OF THE SOUTHWEST, A DOMINANT TECTONIC FORCE THAT WOULD CONTINUE TO SHAPE THE TOPOGRAPHY. THE NEOGENE WAS ALSO A TIME OF HEIGHTENED VOLCANIC ACTIVITY IN MANY PARTS OF THE REGION, CONTRIBUTING SIGNIFICANTLY TO THE GEOLOGICAL MAKEUP. UNDERSTANDING THE DEPOSITIONAL PATTERNS AND ROCK TYPES FROM BOTH THE PALEOGENE AND NEOGENE IS CRUCIAL FOR PIECING TOGETHER THE EVOLUTIONARY NARRATIVE OF THE SOUTHWEST'S GEOLOGICAL PAST, FROM ITS EARLIEST CENOZOIC FORMATIONS TO THE MORE RECENT VOLCANIC FIELDS AND BASIN FILLS.

TECTONIC DRIVERS OF CENOZOIC SOUTHWEST GEOLOGY

THE CENOZOIC GEOLOGICAL HISTORY OF THE SOUTHWEST USA IS INEXTRICABLY LINKED TO POWERFUL TECTONIC FORCES THAT HAVE REPEATEDLY RESHAPED THE EARTH'S CRUST. THE MOST INFLUENTIAL OF THESE FORCES HAS BEEN THE ONGOING INTERACTION BETWEEN THE NORTH AMERICAN PLATE AND THE PACIFIC PLATE. THIS COMPLEX PLATE BOUNDARY HAS RESULTED IN A VARIETY OF DEFORMATION STYLES, INCLUDING COMPRESSION, EXTENSION, AND STRIKE-SLIP FAULTING, EACH LEAVING ITS DISTINCT GEOLOGICAL SIGNATURE ACROSS THE REGION.

BASIN AND RANGE EXTENSION

PERHAPS THE MOST DEFINING TECTONIC PROCESS OF THE CENOZOIC IN THE SOUTHWEST IS THE BASIN AND RANGE EXTENSIONAL

TECTONICS. BEGINNING PRIMARILY IN THE MIOCENE EPOCH OF THE NEOGENE, THIS PROCESS INVOLVES THE STRETCHING AND THINNING OF THE EARTH'S CRUST, LEADING TO THE FORMATION OF NUMEROUS PARALLEL MOUNTAIN RANGES (HORSTS) SEPARATED BY DOWN-DROPPED VALLEYS (GRABENS). THIS TECTONIC REGIME IS RESPONSIBLE FOR THE CHARACTERISTIC, HIGHLY DISSECTED TOPOGRAPHY OF STATES LIKE NEVADA, UTAH, ARIZONA, AND NEW MEXICO. THE STRETCHING OF THE CRUST IS ACCOMPANIED BY SIGNIFICANT FAULTING, WITH LARGE BLOCKS OF ROCK MOVING RELATIVE TO EACH OTHER. THIS ONGOING EXTENSION CONTINUES TO INFLUENCE SEISMIC ACTIVITY IN THE REGION.

COLORADO PLATEAU UPLIFT AND SUBSIDENCE

WHILE MUCH OF THE SOUTHWEST EXPERIENCED EXTENSIONAL FORCES, THE COLORADO PLATEAU, ENCOMPASSING LARGE PARTS OF UTAH, ARIZONA, COLORADO, AND NEW MEXICO, PRESENTS A CONTRASTING GEOLOGICAL STORY. THIS REGION REMAINED RELATIVELY STABLE AND UPLIFTED DURING MUCH OF THE CENOZOIC, SPARED FROM THE INTENSE DEFORMATION SEEN IN ADJACENT AREAS. HOWEVER, EVEN THE COLORADO PLATEAU WAS NOT IMMUNE TO TECTONIC INFLUENCES. DURING THE PALEOGENE, THE LARAMIDE OROGENY CAUSED SIGNIFICANT UPLIFT AND THE FORMATION OF BROAD STRUCTURAL ARCHES AND BASINS WITHIN THE PLATEAU. LATER CENOZOIC PROCESSES, INCLUDING ISOSTATIC ADJUSTMENTS AND REGIONAL TILTING, HAVE ALSO PLAYED A ROLE IN SHAPING ITS ELEVATED LANDSCAPE, SETTING THE STAGE FOR THE IMMENSE EROSIONAL FORCES THAT WOULD LATER CARVE ITS ICONIC CANYONS.

SAN ANDREAS FAULT SYSTEM INFLUENCE

TO THE WEST, THE IMMENSE SAN ANDREAS FAULT SYSTEM, A TRANSFORM FAULT MARKING THE BOUNDARY BETWEEN THE PACIFIC AND NORTH AMERICAN PLATES, EXERTS A SIGNIFICANT INFLUENCE ON THE BROADER TECTONIC SETTING OF THE SOUTHWEST. WHILE NOT DIRECTLY WITHIN THE CORE SOUTHWEST, ITS ACTIVITY HAS LED TO CRUSTAL FRAGMENTATION, THE ROTATION OF TECTONIC BLOCKS, AND THE GENERATION OF STRESSES THAT PROPAGATE INLAND, CONTRIBUTING INDIRECTLY TO SOME OF THE DEFORMATION PATTERNS OBSERVED IN REGIONS LIKE SOUTHERN CALIFORNIA AND WESTERN ARIZONA. THE TRANSFORM MOTION ALONG THE SAN ANDREAS IS A MAJOR DRIVER OF CRUSTAL MOVEMENT AND STRAIN ACCUMULATION ACROSS THE WESTERN CONTINENTAL MARGIN.

VOLCANIC PROVINCES AND THEIR ENDURING LEGACY

THE CENOZOIC WAS AN ERA OF WIDESPREAD AND OFTEN SPECTACULAR VOLCANIC ACTIVITY ACROSS THE SOUTHWEST USA. TECTONIC FORCES, PARTICULARLY THOSE RELATED TO PLATE BOUNDARIES AND CRUSTAL EXTENSION, PROVIDED THE PLUMBING FOR MAGMA TO REACH THE SURFACE, CREATING DIVERSE VOLCANIC LANDSCAPES. THESE ERUPTIONS, RANGING FROM EFFUSIVE LAVA FLOWS TO EXPLOSIVE PYROCLASTIC EVENTS, HAVE LEFT AN INDELIBLE MARK ON THE REGION'S GEOLOGY AND TOPOGRAPHY.

THE SAN JUAN VOLCANIC FIELD

LOCATED PRIMARILY IN SOUTHWESTERN COLORADO AND EXTENDING INTO NORTHERN NEW MEXICO, THE SAN JUAN VOLCANIC FIELD IS ONE OF THE LARGEST AND MOST COMPLEX VOLCANIC PROVINCES IN THE UNITED STATES. MUCH OF ITS ACTIVITY OCCURRED DURING THE OLIGOCENE AND MIOCENE EPOCHS OF THE PALEOGENE AND NEOGENE. MASSIVE ERUPTIONS PRODUCED ENORMOUS VOLUMES OF ASH AND PYROCLASTIC FLOWS, CREATING EXTENSIVE CALDERA SYSTEMS AND THICK IGNIMBRITE SHEETS THAT NOW FORM PROMINENT GEOLOGICAL FEATURES. THE EROSION OF THESE VOLCANIC ROCKS HAS RESULTED IN THE RUGGED, MOUNTAINOUS TERRAIN CHARACTERISTIC OF THE SAN JUAN MOUNTAINS, WITH THEIR DISTINCTIVE JAGGED PEAKS AND DEEP VALLEYS.

THE DATIL-PIE TOWN VOLCANIC FIELD

IN WEST-CENTRAL NEW MEXICO, THE DATIL-PIE TOWN VOLCANIC FIELD REPRESENTS A SIGNIFICANT AREA OF NEOGENE VOLCANIC ACTIVITY. THIS FIELD IS CHARACTERIZED BY NUMEROUS BASALTIC LAVA FLOWS, CINDER CONES, AND SHIELD VOLCANOES. THE RELATIVELY YOUNG AGE OF SOME OF THESE FEATURES, DATING BACK TO THE PLIOCENE AND PLEISTOCENE EPOCHS, INDICATES ONGOING VOLCANIC PROCESSES WITHIN THE REGION. THE DARK, OFTEN VESICULAR BASALTIC ROCKS CONTRAST SHARPLY WITH THE SURROUNDING SEDIMENTARY ROCKS, CREATING DISTINCTIVE GEOLOGICAL FORMATIONS AND INFLUENCING LOCAL SOIL TYPES AND VEGETATION PATTERNS.

OTHER VOLCANIC CENTERS

BEYOND THESE MAJOR FIELDS, NUMEROUS SMALLER VOLCANIC CENTERS AND ISOLATED VOLCANIC FEATURES ARE SCATTERED THROUGHOUT THE CENOZOIC SOUTHWEST. THESE INCLUDE THE JEMEZ MOUNTAINS IN NEW MEXICO, HOME TO THE VALLES CALDERA, A SUPERVOLCANIC CALDERA FORMED BY SEVERAL MASSIVE ERUPTIONS DURING THE PLIOCENE AND PLEISTOCENE. THE MOGOLLON-DATIL VOLCANIC FIELD IN SOUTHWESTERN NEW MEXICO AND SOUTHEASTERN ARIZONA IS ANOTHER EXTENSIVE AREA OF VOLCANIC ACTIVITY, PRIMARILY FEATURING STRATOVOLCANOES AND LAVA FLOWS. THESE DIVERSE VOLCANIC PROVINCES DEMONSTRATE THE WIDESPREAD NATURE OF CENOZOIC MAGMATISM ACROSS THE SOUTHWEST.

SEDIMENTARY BASINS AND DEPOSITIONAL ENVIRONMENTS

INTERSPERSED WITH PERIODS OF MOUNTAIN BUILDING AND VOLCANIC ERUPTIONS, THE CENOZOIC SOUTHWEST WAS ALSO A REALM OF EXTENSIVE SEDIMENT ACCUMULATION IN VAST BASINS. THESE BASINS, OFTEN FORMED BY TECTONIC SUBSIDENCE OR EXTENSIONAL FAULTING, BECAME RECIPIENTS OF ERODED MATERIAL FROM SURROUNDING HIGHLANDS, PRESERVING A DETAILED RECORD OF THE REGION'S ANCIENT ENVIRONMENTS AND CLIMATES.

THE RIO GRANDE RIFT BASINS

A DEFINING FEATURE OF NEW MEXICO'S CENOZOIC GEOLOGY IS THE RIO GRANDE RIFT, A LARGE EXTENSIONAL FEATURE THAT HAS BEEN ACTIVE SINCE THE LATE OLIGOCENE. THIS RIFT IS CHARACTERIZED BY A SERIES OF INTERCONNECTED BASINS, SUCH AS THE ESPANOLA BASIN, THE ALBUQUERQUE BASIN, AND THE TULAROSA BASIN. THESE BASINS HAVE FILLED WITH THICK SEQUENCES OF CENOZOIC SEDIMENTS, INCLUDING ALLUVIAL FAN DEPOSITS, FLUVIAL SANDS AND GRAVELS, AND PLAYA LAKE DEPOSITS. THE INTERPLAY BETWEEN TECTONIC SUBSIDENCE AND SEDIMENT INFILL HAS CREATED DIVERSE DEPOSITIONAL ENVIRONMENTS, PRESERVING FOSSILS AND PROVIDING CRUCIAL INSIGHTS INTO THE PALEOCLIMATE AND PALEOECOLOGY OF THE REGION. THESE BASINS ARE ALSO IMPORTANT SOURCES OF GROUNDWATER AND MINERAL RESOURCES.

THE GREAT BASIN SEDIMENTARY RECORD

THE EXPANSIVE GREAT BASIN, A REGION OF INTERNAL DRAINAGE ENCOMPASSING MUCH OF NEVADA AND PARTS OF UTAH, IS A PRIME EXAMPLE OF A REGION DOMINATED BY CENOZOIC BASIN AND RANGE TOPOGRAPHY AND ITS ASSOCIATED SEDIMENTARY FILL. THOUSANDS OF FEET OF CENOZOIC SEDIMENTS HAVE ACCUMULATED IN THE NUMEROUS BASINS THAT PUNCTUATE THE LANDSCAPE. THESE DEPOSITS INCLUDE VAST ANCIENT LAKEBEDS, SUCH AS THOSE OF PREHISTORIC LAKE BONNEVILLE, WHICH LEFT BEHIND EXTENSIVE SHORELINES AND LACUSTRINE SEDIMENTS. THE SEDIMENTARY RECORD HERE IS CRUCIAL FOR UNDERSTANDING PALEOHYDROLOGY, CLIMATE CHANGE, AND THE EVOLUTION OF THE UNIQUE GREAT BASIN ECOSYSTEM.

FLUVIAL AND ALLUVIAL DEPOSITS

ACROSS MUCH OF THE CENOZOIC SOUTHWEST, FLUVIAL (RIVER) AND ALLUVIAL (ANCIENT RIVER AND FAN) DEPOSITS ARE UBIQUITOUS. THESE SEDIMENTS, CONSISTING OF SANDS, GRAVELS, AND MUDS, WERE TRANSPORTED BY RIVERS DRAINING THE UPLIFTED MOUNTAIN RANGES AND DEPOSITED IN LOW-LYING AREAS. FORMATIONS LIKE THE SALT WASH MEMBER OF THE MORRISON FORMATION (THOUGH MESOZOIC, ITS CENOZOIC DEPOSITIONAL CONTINUATION IS RELEVANT IN UNDERSTANDING LATER ALLUVIAL FAN DEPOSITS) AND VARIOUS NEOGENE TERRESTRIAL FORMATIONS ARE TESTAMENT TO THE VAST RIVER SYSTEMS AND EXTENSIVE FLOODPLAINS THAT ONCE EXISTED. THESE DEPOSITS ARE OFTEN RICH IN FOSSILS AND REPRESENT CRUCIAL SOURCES OF BUILDING MATERIALS AND AGGREGATE.

EROSIONAL SCULPTURES: THE ARID SOUTHWEST'S MASTERPIECES

THE RELATIVE ARIDITY OF THE SOUTHWEST USA, ESPECIALLY DURING THE LATER CENOZOIC, HAS BEEN A CRITICAL FACTOR IN SHAPING ITS ICONIC LANDSCAPES THROUGH EROSIONAL PROCESSES. ONCE UPLIFTED AND EXPOSED, THE DIVERSE GEOLOGICAL MATERIALS OF THE CENOZOIC HAVE BEEN SUBJECT TO THE RELENTLESS WORK OF WATER, WIND, AND ICE, CARVING INTRICATE AND AWE-INSPIRING LANDFORMS.

CANYON FORMATION AND MESA TOPOGRAPHY

THE GRAND CANYON, A TESTAMENT TO MILLIONS OF YEARS OF EROSION INTO THE UPLIFTED COLORADO PLATEAU, IS THE ULTIMATE CENOZOIC MASTERPIECE SCULPTED BY THE COLORADO RIVER. HOWEVER, NUMEROUS OTHER CANYONS, ARROYOS, AND INCISED RIVER VALLEYS CRISSCROSS THE SOUTHWEST, CARVED INTO CENOZOIC SEDIMENTARY AND VOLCANIC ROCKS. THE DIFFERENTIAL EROSION OF THESE MATERIALS, WHERE HARDER LAYERS RESIST WEATHERING WHILE SOFTER ONES ARE WORN AWAY, LEADS TO THE FORMATION OF MESAS, BUTTES, AND SPIRES. THE ARID CLIMATE LIMITS THE VEGETATION COVER, ALLOWING FOR MORE EFFICIENT EROSION AND THE PRESERVATION OF THESE DRAMATIC LANDFORMS.

WIND EROSION AND DESERT PAVEMENT

WIND EROSION, OR AEOLIAN PROCESSES, PLAYS A SIGNIFICANT ROLE IN SHAPING CENOZOIC LANDSCAPES IN THE ARID SOUTHWEST. DUST AND SAND ARE TRANSPORTED AND DEPOSITED, CREATING FEATURES LIKE SAND DUNES AND ERGS. IN MANY AREAS, THE REMOVAL OF FINER PARTICLES BY WIND LEAVES BEHIND A SURFACE LAYER OF LARGER PEBBLES AND ROCKS, KNOWN AS DESERT PAVEMENT. THIS PROCESS IS EVIDENT ACROSS VAST EXPANSES OF THE REGION, CONTRIBUTING TO THE CHARACTERISTIC STARK BEAUTY OF DESERT ENVIRONMENTS. THE FINE DUST DEPOSITED FROM WIND CAN ALSO CONTRIBUTE TO THE FORMATION OF FERTILE SOILS IN AREAS WHERE IT ACCUMULATES.

FORMATION OF ARCHES AND HOODOOS

THE UNIQUE GEOLOGICAL FORMATIONS OF ARCHES AND HOODOOS, OFTEN FOUND IN SANDSTONE FORMATIONS OF CENOZOIC AGE, ARE A DIRECT RESULT OF WEATHERING AND EROSION. DIFFERENTIAL EROSION, WHERE SOFTER ROCK IS REMOVED MORE QUICKLY THAN HARDER ROCK, CREATES THESE FANTASTICAL SHAPES. WATER PERCOLATING THROUGH JOINTS AND CRACKS IN THE ROCK, COUPLED WITH FREEZE-THAW CYCLES AND WIND ABRASION, SCULPTS THESE ICONIC NATURAL SCULPTURES. EXAMPLES CAN BE FOUND IN NUMEROUS NATIONAL AND STATE PARKS THROUGHOUT THE SOUTHWEST, SHOWCASING THE CREATIVE POWER OF EROSIONAL FORCES ACTING ON CENOZOIC ROCK UNITS.

SIGNIFICANT CENOZOIC GEOLOGICAL FEATURES OF THE SOUTHWEST

THE CENOZOIC GEOLOGICAL HISTORY OF THE SOUTHWEST USA HAS RESULTED IN A STUNNING ARRAY OF DISTINCTIVE FEATURES THAT CONTINUE TO FASCINATE GEOLOGISTS AND VISITORS ALIKE. THESE FEATURES ARE DIRECT PRODUCTS OF THE TECTONIC, VOLCANIC, AND EROSIONAL PROCESSES DISCUSSED PREVIOUSLY.

- **THE GRAND CANYON:** A COLOSSAL TESTAMENT TO THE EROSIONAL POWER OF THE COLORADO RIVER ACTING ON UPLIFTED PALEOZOIC AND MESOZOIC STRATA, BUT ITS FORMATION AND PRESERVATION ARE INTRICATELY LINKED TO THE CENOZOIC UPLIFT AND ARID CLIMATE.
- **MONUMENT VALLEY:** FAMOUS FOR ITS TOWERING SANDSTONE MESAS AND BUTTES, SCULPTED BY EROSION FROM SEDIMENTARY LAYERS DEPOSITED DURING THE MESOZOIC AND LATER SCULPTED BY CENOZOIC WEATHERING.
- **THE SUPERSTITION MOUNTAINS:** A RUGGED RANGE IN ARIZONA, CHARACTERIZED BY VOLCANIC ROCKS FROM CENOZOIC ERUPTIONS, INCLUDING IGNIMBRITES AND ANDESITIC LAVA FLOWS, SCULPTED INTO DRAMATIC PEAKS AND CANYONS.
- **DEATH VALLEY NATIONAL PARK:** A PRIME EXAMPLE OF BASIN AND RANGE TOPOGRAPHY, FEATURING DEEP GRABENS FILLED WITH CENOZOIC SEDIMENTS AND ANCIENT LAKEBEDS, ADJACENT TO UPLIFTED FAULT BLOCKS.
- **THE PAINTED DESERT:** A VAST EXPANSE OF COLORFUL BADLANDS, PREDOMINANTLY COMPOSED OF TERTIARY (PALEOGENE AND NEOGENE) SEDIMENTARY ROCKS THAT HAVE BEEN INTRICATELY ERODED BY WIND AND WATER INTO A SURREAL LANDSCAPE.
- **VARIOUS CALDERAS AND VOLCANIC NECKS:** NUMEROUS FEATURES ACROSS ARIZONA, NEW MEXICO, AND COLORADO, SUCH AS THE VALLES CALDERA AND SHIPROCK, REPRESENT THE DRAMATIC VOLCANIC LEGACY OF THE CENOZOIC ERA.

THESE FEATURES, AND MANY OTHERS, ARE THE TANGIBLE RESULTS OF MILLIONS OF YEARS OF DYNAMIC GEOLOGICAL PROCESSES. THEIR STUDY PROVIDES INVALUABLE INSIGHTS INTO THE EARTH'S HISTORY AND THE ONGOING EVOLUTION OF OUR PLANET. THE CENOZOIC ERA IN THE SOUTHWEST IS A CONTINUING STORY, WITH TECTONIC ACTIVITY, VOLCANIC POTENTIAL, AND EROSIONAL FORCES STILL ACTIVELY SHAPING THE LAND.

FAQ

Q: WHAT IS THE MOST SIGNIFICANT TECTONIC EVENT THAT SHAPED CENOZOIC GEOLOGY IN THE SOUTHWEST USA?

A: THE MOST SIGNIFICANT TECTONIC EVENT IS THE BASIN AND RANGE EXTENSION, WHICH BEGAN PRIMARILY IN THE MIOCENE EPOCH AND CONTINUES TODAY. THIS PROCESS INVOLVES THE STRETCHING AND THINNING OF THE EARTH'S CRUST, LEADING TO THE CHARACTERISTIC TOPOGRAPHY OF PARALLEL MOUNTAIN RANGES (HORSTS) AND DOWN-DROPPED VALLEYS (GRABENS) ACROSS MUCH OF THE SOUTHWEST.

Q: WHAT TYPES OF VOLCANIC ROCKS ARE MOST COMMON IN THE CENOZOIC SOUTHWEST?

A: THE MOST COMMON VOLCANIC ROCKS INCLUDE BASALTIC LAVAS, ANDESITES, RHYOLITES, AND EXTENSIVE DEPOSITS OF VOLCANIC ASH AND PYROCLASTIC FLOWS (IGNIMBRITES). THESE ROCKS ARE FOUND IN LARGE VOLCANIC FIELDS, CALDERAS, AND AS ISOLATED VOLCANIC CONES AND LAVA FLOWS.

Q: HOW DID THE ARID CLIMATE INFLUENCE CENOZOIC GEOLOGY IN THE SOUTHWEST?

A: THE ARID CLIMATE SIGNIFICANTLY INFLUENCED CENOZOIC GEOLOGY BY PROMOTING EROSIONAL PROCESSES LIKE WIND (AEOLIAN) EROSION AND THE RAPID CARVING OF CANYONS AND VALLEYS BY WATER. LIMITED VEGETATION COVER IN ARID REGIONS ALLOWS FOR MORE DIRECT AND EFFICIENT WEATHERING AND EROSION OF ROCK FORMATIONS, LEADING TO THE FORMATION OF DRAMATIC LANDFORMS LIKE MESAS, BUTTES, AND ARCHES.

Q: WHAT ARE SOME OF THE KEY SEDIMENTARY BASINS FORMED DURING THE CENOZOIC IN THE SOUTHWEST USA?

A: PROMINENT CENOZOIC SEDIMENTARY BASINS INCLUDE THE RIO GRANDE RIFT BASINS (E.G., ESPANOLA BASIN, ALBUQUERQUE BASIN), THE BASINS OF THE GREAT BASIN REGION (E.G., LAKE BONNEVILLE BASIN), AND VARIOUS SMALLER INTERMONTANE BASINS FORMED BY BASIN AND RANGE EXTENSION. THESE BASINS ARE FILLED WITH THICK SEQUENCES OF TERRESTRIAL SEDIMENTS.

Q: CAN YOU GIVE AN EXAMPLE OF A MAJOR CENOZOIC CALDERA IN THE SOUTHWEST?

A: THE VALLES CALDERA IN THE JEMEZ MOUNTAINS OF NEW MEXICO IS A SIGNIFICANT EXAMPLE OF A CENOZOIC CALDERA. IT WAS FORMED BY MASSIVE SUPERVOLCANIC ERUPTIONS DURING THE PLIOCENE AND PLEISTOCENE EPOCHS AND IS A HIGHLY STUDIED FEATURE DUE TO ITS GEOLOGICAL IMPORTANCE AND POTENTIAL FOR GEOTHERMAL ENERGY.

Q: WHAT ROLE DID THE LARAMIDE OROGENY PLAY IN CENOZOIC SOUTHWEST GEOLOGY?

A: WHILE THE LARAMIDE OROGENY'S MOST INTENSE MOUNTAIN-BUILDING OCCURRED AT THE END OF THE MESOZOIC AND THE VERY BEGINNING OF THE CENOZOIC, ITS COMPRESSIONAL FORCES CONTINUED TO INFLUENCE CRUSTAL DEFORMATION IN THE SOUTHWEST DURING THE PALEOGENE. IT CAUSED SIGNIFICANT UPLIFT AND THE FORMATION OF BROAD STRUCTURAL ARCHES AND BASINS, PARTICULARLY ON THE COLORADO PLATEAU, SETTING THE STAGE FOR SUBSEQUENT CENOZOIC GEOLOGICAL EVENTS.

Q: HOW DOES CENOZOIC GEOLOGY CONTRIBUTE TO THE WATER RESOURCES OF THE SOUTHWEST?

A: CENOZOIC SEDIMENTARY BASINS, PARTICULARLY THOSE ASSOCIATED WITH THE RIO GRANDE RIFT AND THE GREAT BASIN, ARE CRITICAL AQUIFERS, STORING VAST AMOUNTS OF GROUNDWATER WITHIN THEIR POROUS AND PERMEABLE SEDIMENT LAYERS. THE GEOLOGICAL STRUCTURE AND COMPOSITION OF THESE BASINS DIRECTLY INFLUENCE GROUNDWATER RECHARGE, FLOW, AND AVAILABILITY, MAKING THEM VITAL FOR HUMAN POPULATIONS AND ECOSYSTEMS IN THE ARID SOUTHWEST.

[Cenozoic Geology Southwest Usa](#)

Cenozoic Geology Southwest Usa

Related Articles

- [challenges to us soft power](#)
- [certified electronics recycler](#)
- [championship sports physics](#)

[Back to Home](#)