

DARK MATTER AND ITS PROPERTIES

THE COSMIC ENIGMA: UNVEILING DARK MATTER AND ITS PROPERTIES

THE INVISIBLE INFLUENCE: WHAT EXACTLY IS DARK MATTER?

DARK MATTER AND ITS PROPERTIES ARE CENTRAL TO OUR UNDERSTANDING OF THE UNIVERSE, YET THEY REMAIN ONE OF COSMOLOGY'S MOST PROFOUND MYSTERIES. WHILE WE CAN OBSERVE AND INTERACT WITH ORDINARY MATTER – THE STARS, PLANETS, GALAXIES, AND OURSELVES – A VAST AMOUNT OF THE UNIVERSE'S MASS APPEARS TO BE ENTIRELY INVISIBLE. THIS ELUSIVE SUBSTANCE, DUBBED "DARK MATTER," DOESN'T EMIT, ABSORB, OR REFLECT LIGHT, MAKING IT UNDETECTABLE BY CONVENTIONAL TELESCOPES. ITS PRESENCE IS INFERRED SOLELY THROUGH ITS GRAVITATIONAL EFFECTS ON VISIBLE MATTER AND LIGHT. SCIENTISTS ESTIMATE THAT DARK MATTER CONSTITUTES ABOUT 27% OF THE UNIVERSE'S TOTAL MASS-ENERGY CONTENT, DWARFING THE MERE 5% THAT MAKES UP THE ORDINARY, BARYONIC MATTER WE ARE FAMILIAR WITH. THE REMAINING 68% IS ATTRIBUTED TO DARK ENERGY, ANOTHER ENIGMATIC FORCE DRIVING THE ACCELERATED EXPANSION OF THE COSMOS. UNDERSTANDING DARK MATTER IS CRUCIAL NOT ONLY FOR COMPREHENDING THE UNIVERSE'S CURRENT STRUCTURE BUT ALSO FOR DECIPHERING ITS PAST EVOLUTION AND PREDICTING ITS ULTIMATE FATE. THIS ARTICLE DELVES INTO THE COMPELLING EVIDENCE FOR DARK MATTER, EXPLORES ITS INFERRED PROPERTIES, AND DISCUSSES THE ONGOING QUESTS TO IDENTIFY ITS FUNDAMENTAL NATURE.

TABLE OF CONTENTS

- THE INVISIBLE INFLUENCE: WHAT EXACTLY IS DARK MATTER?
- UNVEILING THE EVIDENCE: WHY WE BELIEVE DARK MATTER EXISTS
- GALACTIC ROTATION CURVES: THE FIRST CLUES
- GRAVITATIONAL LENSING: BENDING LIGHT, REVEALING MASS
- COSMIC MICROWAVE BACKGROUND: ECHOES OF THE EARLY UNIVERSE
- GALAXY CLUSTERS: THE MISSING MASS PUZZLE
- THE INFERRED PROPERTIES OF DARK MATTER
- GRAVITATIONAL INTERACTION: ITS DEFINING CHARACTERISTIC
- NON-BARYONIC NATURE: NOT MADE OF PROTONS AND NEUTRONS
- WEAK INTERACTION: THE ELUSIVE NATURE OF ITS OTHER FORCES
- COLD VS. HOT DARK MATTER: MODELS AND PREDICTIONS
- DARK MATTER CANDIDATES: THE SEARCH FOR THE UNSEEN PARTICLE
- WIMPS: WEAKLY INTERACTING MASSIVE PARTICLES
- AXIONS: ULTRA-LIGHT PARTICLES WITH UNIQUE PROPERTIES
- MACHOS: MASSIVE ASTROPHYSICAL COMPACT HALO OBJECTS
- STERILE NEUTRINOS: A HYPOTHETICAL DARK MATTER PARTICLE

- DIRECT DETECTION EXPERIMENTS: LISTENING FOR THE WHISPER
- INDIRECT DETECTION EXPERIMENTS: HUNTING FOR COSMIC BYPRODUCTS
- COLLIDER EXPERIMENTS: FORGING DARK MATTER IN THE LAB
- THE ONGOING QUEST AND FUTURE PROSPECTS

UNVEILING THE EVIDENCE: WHY WE BELIEVE DARK MATTER EXISTS

THE NOTION OF DARK MATTER WASN'T BORN FROM A WILD GUESS; IT EMERGED FROM CAREFUL ASTRONOMICAL OBSERVATIONS THAT COULDN'T BE EXPLAINED BY THE VISIBLE MATTER ALONE. FOR DECADES, DISCREPANCIES BETWEEN THEORETICAL PREDICTIONS BASED ON OBSERVABLE MASS AND ACTUAL OBSERVED PHENOMENA HAVE ACCUMULATED, PAINTING A CONSISTENT PICTURE OF AN UNSEEN GRAVITATIONAL ANCHOR HOLDING THE COSMOS TOGETHER. THESE PIECES OF EVIDENCE, GATHERED FROM VARIOUS COSMOLOGICAL SCALES, FORM A COMPELLING CASE FOR THE EXISTENCE OF SOMETHING BEYOND ORDINARY MATTER.

GALACTIC ROTATION CURVES: THE FIRST CLUES

ONE OF THE EARLIEST AND MOST PERSUASIVE PIECES OF EVIDENCE FOR DARK MATTER CAME FROM STUDYING HOW GALAXIES ROTATE. IN THE LATE 1970S, VERA RUBIN AND KENT FORD METICULOUSLY OBSERVED THE SPEEDS AT WHICH STARS ORBIT THE CENTERS OF SPIRAL GALAXIES. ACCORDING TO NEWTONIAN PHYSICS, STARS FARTHER AWAY FROM THE GALACTIC CENTER, WHERE MOST OF THE VISIBLE MASS IS CONCENTRATED, SHOULD ORBIT MORE SLOWLY, MUCH LIKE PLANETS FARTHER FROM THE SUN MOVE AT LOWER SPEEDS. HOWEVER, RUBIN AND FORD DISCOVERED SOMETHING ASTONISHING: STARS ON THE OUTER EDGES OF GALAXIES WERE ORBITING JUST AS FAST, IF NOT FASTER, THAN THOSE CLOSER TO THE CENTER. THIS IMPLIED THAT THERE WAS A SIGNIFICANT AMOUNT OF UNSEEN MASS DISTRIBUTED THROUGHOUT THE GALAXY, EXTENDING FAR BEYOND THE VISIBLE DISC AND BULGE, PROVIDING THE EXTRA GRAVITATIONAL PULL NEEDED TO KEEP THESE FAST-MOVING STARS IN THEIR ORBITS. THIS UNSEEN MASS IS WHAT WE NOW REFER TO AS THE GALACTIC HALO, AND IT IS PREDOMINANTLY COMPOSED OF DARK MATTER.

GRAVITATIONAL LENSING: BENDING LIGHT, REVEALING MASS

ANOTHER POWERFUL INDICATOR OF DARK MATTER'S PRESENCE IS GRAVITATIONAL LENSING. ALBERT EINSTEIN'S THEORY OF GENERAL RELATIVITY PREDICTS THAT MASSIVE OBJECTS WARP THE FABRIC OF SPACETIME, CAUSING LIGHT TO BEND AS IT PASSES BY. THIS PHENOMENON IS AKIN TO A LENS BENDING LIGHT. ASTRONOMERS OBSERVE THAT LIGHT FROM DISTANT GALAXIES IS OFTEN DISTORTED, MAGNIFIED, OR EVEN APPEARS AS MULTIPLE IMAGES WHEN IT TRAVELS THROUGH REGIONS OF SPACE CONTAINING LARGE CONCENTRATIONS OF MASS, SUCH AS GALAXY CLUSTERS. BY ANALYZING THE DEGREE OF LIGHT BENDING, SCIENTISTS CAN MAP THE DISTRIBUTION OF MASS IN THESE REGIONS. CONSISTENTLY, THESE LENSING STUDIES REVEAL THAT THE TOTAL MASS INFERRED FROM THE BENDING OF LIGHT IS SIGNIFICANTLY GREATER THAN THE MASS ACCOUNTED FOR BY THE VISIBLE STARS AND GAS. THIS "MISSING MASS" ALIGNS PERFECTLY WITH THE DISTRIBUTION EXPECTED FOR DARK MATTER, ACTING AS AN INVISIBLE GRAVITATIONAL LENS.

COSMIC MICROWAVE BACKGROUND: ECHOES OF THE EARLY UNIVERSE

THE COSMIC MICROWAVE BACKGROUND (CMB) RADIATION IS THE FAINT AFTERGLOW OF THE BIG BANG, A SNAPSHOT OF THE UNIVERSE WHEN IT WAS JUST ABOUT 380,000 YEARS OLD. STUDYING THE SUBTLE TEMPERATURE FLUCTUATIONS IN THE CMB HAS PROVIDED CRUCIAL INSIGHTS INTO THE EARLY UNIVERSE'S COMPOSITION AND EVOLUTION. COSMOLOGICAL MODELS THAT INCORPORATE DARK MATTER ARE REMARKABLY SUCCESSFUL AT EXPLAINING THE OBSERVED PATTERNS OF THESE FLUCTUATIONS.

SPECIFICALLY, THE RELATIVE HEIGHTS OF THE PEAKS IN THE CMB'S POWER SPECTRUM STRONGLY SUGGEST THAT THE UNIVERSE CONTAINS A SUBSTANTIAL AMOUNT OF NON-BARYONIC DARK MATTER, WHICH PLAYED A CRUCIAL ROLE IN SEEDING THE FORMATION OF THE LARGE-SCALE STRUCTURES WE SEE TODAY, SUCH AS GALAXIES AND CLUSTERS.

GALAXY CLUSTERS: THE MISSING MASS PUZZLE

GALAXY CLUSTERS ARE THE LARGEST GRAVITATIONALLY BOUND STRUCTURES IN THE UNIVERSE, CONTAINING HUNDREDS OR EVEN THOUSANDS OF GALAXIES. OBSERVATIONS OF THESE COLOSSAL STRUCTURES HAVE ALSO REVEALED A STARK DISCREPANCY BETWEEN VISIBLE AND TOTAL MASS. GALAXIES WITHIN CLUSTERS MOVE AT VERY HIGH SPEEDS, AND IF ONLY THE VISIBLE MATTER WERE PRESENT, THESE CLUSTERS WOULD HAVE DISPERSED LONG AGO. THE FACT THAT THEY REMAIN BOUND INDICATES A FAR GREATER GRAVITATIONAL PULL. FURTHERMORE, THE HOT GAS OBSERVED WITHIN GALAXY CLUSTERS, WHICH EMITS X-RAYS, IS MUCH HOTTER THAN EXPECTED. THIS IMPLIES THAT THE GAS IS CONFINED BY A MUCH STRONGER GRAVITATIONAL FIELD THAN CAN BE ACCOUNTED FOR BY THE VISIBLE GALAXIES ALONE. BOTH THE DYNAMICS OF GALAXIES AND THE CONFINEMENT OF HOT GAS POINT TO THE OVERWHELMING DOMINANCE OF DARK MATTER IN HOLDING THESE MASSIVE STRUCTURES TOGETHER.

THE INFERRED PROPERTIES OF DARK MATTER

WHILE WE CANNOT DIRECTLY SEE DARK MATTER, ITS OBSERVED GRAVITATIONAL EFFECTS ALLOW US TO INFER SEVERAL KEY PROPERTIES. THESE PROPERTIES ARE CRUCIAL FOR GUIDING THE SEARCH FOR ITS FUNDAMENTAL NATURE AND FOR DEVELOPING THEORETICAL MODELS THAT CAN EXPLAIN ITS EXISTENCE.

GRAVITATIONAL INTERACTION: ITS DEFINING CHARACTERISTIC

THE MOST FUNDAMENTAL AND DEFINING PROPERTY OF DARK MATTER IS ITS INTERACTION THROUGH GRAVITY. IT EXERTS A GRAVITATIONAL PULL ON ORDINARY MATTER AND ON ITSELF, SHAPING THE LARGE-SCALE STRUCTURE OF THE UNIVERSE, FROM THE FORMATION OF GALAXIES TO THE CLUSTERING OF GALAXY CLUSTERS. WITHOUT THIS GRAVITATIONAL INFLUENCE, THE UNIVERSE WOULD LOOK VASTLY DIFFERENT, AND THE STRUCTURES WE OBSERVE WOULD NOT HAVE HAD TIME TO FORM. THIS GRAVITATIONAL INTERACTION IS THE SOLE REASON WE KNOW DARK MATTER EXISTS.

NON-BARYONIC NATURE: NOT MADE OF PROTONS AND NEUTRONS

ONE OF THE MOST SIGNIFICANT INFERRED PROPERTIES IS THAT DARK MATTER IS NOT COMPOSED OF ORDINARY BARYONIC MATTER – PROTONS, NEUTRONS, AND ELECTRONS. THIS CONCLUSION STEMS FROM SEVERAL LINES OF EVIDENCE, INCLUDING THE ABUNDANCE OF LIGHT ELEMENTS FORGED IN THE BIG BANG (BIG BANG NUCLEOSYNTHESIS) AND THE PATTERNS OBSERVED IN THE COSMIC MICROWAVE BACKGROUND. THESE OBSERVATIONS CONSTRAIN THE TOTAL AMOUNT OF BARYONIC MATTER IN THE UNIVERSE TO A MUCH LOWER LEVEL THAN THE TOTAL MASS INFERRED FROM GRAVITATIONAL EFFECTS. THEREFORE, DARK MATTER MUST BE MADE OF SOMETHING ELSE, SOMETHING FUNDAMENTALLY DIFFERENT FROM THE MATTER THAT CONSTITUTES STARS, PLANETS, AND US.

WEAK INTERACTION: THE ELUSIVE NATURE OF ITS OTHER FORCES

BEYOND GRAVITY, DARK MATTER APPEARS TO INTERACT VERY WEAKLY, IF AT ALL, WITH ELECTROMAGNETIC FORCES AND POSSIBLY THE STRONG NUCLEAR FORCE. THE FACT THAT IT DOESN'T EMIT OR ABSORB LIGHT (ELECTROMAGNETIC FORCE) IS ITS NAMESAKE CHARACTERISTIC – IT'S "DARK." THIS LACK OF STRONG INTERACTION WITH LIGHT MEANS IT DOESN'T CLUMP TOGETHER IN THE SAME WAY ORDINARY MATTER DOES TO FORM STARS AND PLANETS. ITS WEAK INTERACTION WITH OTHER FORCES ALSO EXPLAINS WHY IT'S SO DIFFICULT TO DETECT DIRECTLY. WHILE SOME THEORETICAL MODELS PROPOSE A WEAK

INTERACTION WITH ORDINARY MATTER (HENCE THE NAME "WEAKLY INTERACTING MASSIVE PARTICLES" OR WIMPS), THIS INTERACTION IS EXTREMELY SUBTLE AND HAS YET TO BE DEFINITELY OBSERVED.

COLD VS. HOT DARK MATTER: MODELS AND PREDICTIONS

THEORIES ABOUT DARK MATTER OFTEN DISTINGUISH BETWEEN "COLD" AND "HOT" DARK MATTER, BASED ON THE SPEED OF ITS PARTICLES IN THE EARLY UNIVERSE. COLD DARK MATTER (CDM) CONSISTS OF SLOW-MOVING PARTICLES THAT WOULD HAVE STARTED CLUMPING TOGETHER EARLY ON, FORMING THE SEEDS FOR THE HIERARCHICAL STRUCTURE FORMATION WE OBSERVE TODAY, WHERE SMALLER STRUCTURES MERGE TO FORM LARGER ONES. HOT DARK MATTER (HDM) WOULD HAVE BEEN RELATIVISTIC (MOVING AT OR NEAR THE SPEED OF LIGHT) IN THE EARLY UNIVERSE, WHICH WOULD HAVE SMOOTHED OUT INITIAL DENSITY FLUCTUATIONS, LEADING TO A TOP-DOWN STRUCTURE FORMATION SCENARIO (LARGE STRUCTURES FORMING FIRST AND THEN FRAGMENTING). CURRENT COSMOLOGICAL OBSERVATIONS STRONGLY FAVOR THE CDM MODEL, AS IT BETTER EXPLAINS THE OBSERVED DISTRIBUTION OF GALAXIES AND GALAXY CLUSTERS. THEREFORE, MOST DARK MATTER CANDIDATES ARE EXPECTED TO BE "COLD" OR "WARM" (AN INTERMEDIATE CATEGORY) RATHER THAN "HOT."

DARK MATTER CANDIDATES: THE SEARCH FOR THE UNSEEN PARTICLE

THE ELUSIVE NATURE OF DARK MATTER HAS SPURRED A VAST THEORETICAL AND EXPERIMENTAL EFFORT TO IDENTIFY ITS CONSTITUENT PARTICLE. NUMEROUS CANDIDATES HAVE BEEN PROPOSED, EACH WITH UNIQUE PROPERTIES AND DETECTION STRATEGIES.

WIMPS: WEAKLY INTERACTING MASSIVE PARTICLES

PERHAPS THE MOST POPULAR CANDIDATES FOR DARK MATTER ARE WEAKLY INTERACTING MASSIVE PARTICLES, OR WIMPS. THESE HYPOTHETICAL PARTICLES ARE PREDICTED BY SOME EXTENSIONS OF THE STANDARD MODEL OF PARTICLE PHYSICS, SUCH AS SUPERSYMMETRY. WIMPS WOULD BE MASSIVE, MEANING THEY HAVE A SIGNIFICANT REST MASS, AND INTERACT ONLY THROUGH GRAVITY AND THE WEAK NUCLEAR FORCE. THEIR PREDICTED PROPERTIES ALIGN WELL WITH THE REQUIREMENTS OF COLD DARK MATTER. EXPERIMENTS AROUND THE WORLD ARE ACTIVELY SEARCHING FOR WIMPS BY LOOKING FOR THE RARE OCCASIONS WHEN A WIMP MIGHT COLLIDE WITH AN ATOMIC NUCLEUS IN A DETECTOR, PRODUCING A FAINT SIGNAL.

AXIONS: ULTRA-LIGHT PARTICLES WITH UNIQUE PROPERTIES

ANOTHER INTRIGUING CANDIDATE IS THE AXION. PROPOSED TO SOLVE A PROBLEM IN QUANTUM CHROMODYNAMICS (QCD), AXIONS ARE PREDICTED TO BE VERY LIGHT PARTICLES WITH EXTREMELY WEAK INTERACTIONS. THEIR ULTRALIGHT NATURE MEANS THAT A VAST NUMBER OF THEM WOULD BE NEEDED TO ACCOUNT FOR THE OBSERVED DARK MATTER DENSITY. THE SEARCH FOR AXIONS INVOLVES LOOKING FOR THEIR POTENTIAL CONVERSION INTO PHOTONS IN THE PRESENCE OF STRONG MAGNETIC FIELDS, A PROCESS THAT REQUIRES HIGHLY SENSITIVE DETECTION EQUIPMENT.

MACHOS: MASSIVE ASTROPHYSICAL COMPACT HALO OBJECTS

MACHOS, OR MASSIVE ASTROPHYSICAL COMPACT HALO OBJECTS, REPRESENT A CLASS OF DARK MATTER CANDIDATES THAT ARE BARYONIC IN NATURE BUT VERY DIFFICULT TO DETECT. THESE COULD INCLUDE BROWN DWARFS, FAINT WHITE DWARFS, NEUTRON STARS, OR EVEN PRIMORDIAL BLACK HOLES THAT DO NOT EMIT MUCH LIGHT. SEARCHES FOR MACHOS HAVE HISTORICALLY INVOLVED LOOKING FOR GRAVITATIONAL MICROLENSING EVENTS, WHERE A MACHO PASSING IN FRONT OF A DISTANT STAR BRIEFLY MAGNIFIES ITS LIGHT. WHILE MICROLENSING SEARCHES HAVE RULED OUT MACHOS AS THE PRIMARY COMPONENT OF DARK MATTER, THEY HAVEN'T ENTIRELY EXCLUDED THEM AS A MINOR CONTRIBUTOR.

STERILE NEUTRINOS: A HYPOTHETICAL DARK MATTER PARTICLE

NEUTRINOS ARE KNOWN PARTICLES THAT HAVE A VERY SMALL MASS AND INTERACT ONLY THROUGH THE WEAK FORCE AND GRAVITY. STANDARD NEUTRINOS ARE TOO LIGHT AND TOO HOT TO BE THE DOMINANT FORM OF DARK MATTER. HOWEVER, THEORETICAL EXTENSIONS TO THE STANDARD MODEL PROPOSE THE EXISTENCE OF "STERILE NEUTRINOS," WHICH WOULD INTERACT EVEN MORE WEAKLY THAN REGULAR NEUTRINOS AND COULD POTENTIALLY HAVE SUFFICIENT MASS TO BE A DARK MATTER CANDIDATE. DETECTING STERILE NEUTRINOS IS INCREDIBLY CHALLENGING DUE TO THEIR EVEN WEAKER INTERACTIONS.

THE ONGOING QUEST AND FUTURE PROSPECTS

THE QUEST TO UNRAVEL THE MYSTERY OF DARK MATTER IS ONE OF THE MOST EXCITING FRONTIERS IN MODERN PHYSICS AND ASTRONOMY. RESEARCHERS ARE EMPLOYING A MULTI-PRONGED APPROACH, COMBINING THEORETICAL ADVANCEMENTS WITH INCREASINGLY SOPHISTICATED EXPERIMENTAL TECHNIQUES. DIRECT DETECTION EXPERIMENTS, BURIED DEEP UNDERGROUND TO SHIELD THEM FROM COSMIC RAYS, ARE METICULOUSLY DESIGNED TO CAPTURE THE MINUSCULE RECOIL ENERGY WHEN A DARK MATTER PARTICLE INTERACTS WITH DETECTOR MATERIAL. INDIRECT DETECTION EXPERIMENTS, OFTEN SPACE-BASED OR AT HIGH ALTITUDES, SEARCH FOR THE ANNIHILATION OR DECAY PRODUCTS OF DARK MATTER PARTICLES, SUCH AS GAMMA RAYS OR NEUTRINOS, THAT MIGHT BE PRODUCED IN DENSE REGIONS LIKE THE GALACTIC CENTER. MEANWHILE, PARTICLE COLLIDERS LIKE THE LARGE HADRON COLLIDER ARE ATTEMPTING TO CREATE DARK MATTER PARTICLES IN HIGH-ENERGY COLLISIONS. THE FUTURE HOLDS THE PROMISE OF NEW, MORE SENSITIVE DETECTORS AND POTENTIALLY GROUNDBREAKING DISCOVERIES THAT COULD FINALLY REVEAL THE FUNDAMENTAL NATURE OF THIS INVISIBLE COSMIC CONSTITUENT, REVOLUTIONIZING OUR UNDERSTANDING OF THE UNIVERSE.

FREQUENTLY ASKED QUESTIONS ABOUT DARK MATTER AND ITS PROPERTIES

Q: WHAT IS THE PRIMARY EVIDENCE FOR THE EXISTENCE OF DARK MATTER?

A: THE PRIMARY EVIDENCE FOR DARK MATTER COMES FROM ITS GRAVITATIONAL EFFECTS, OBSERVED THROUGH GALACTIC ROTATION CURVES, GRAVITATIONAL LENSING AROUND GALAXY CLUSTERS, THE PATTERNS IN THE COSMIC MICROWAVE BACKGROUND RADIATION, AND THE DYNAMICS OF GALAXY CLUSTERS THEMSELVES. THESE OBSERVATIONS CONSISTENTLY SHOW MORE MASS THAN CAN BE ACCOUNTED FOR BY VISIBLE MATTER.

Q: HOW DOES DARK MATTER INTERACT WITH ORDINARY MATTER?

A: DARK MATTER INTERACTS PRIMARILY THROUGH GRAVITY. IT DOES NOT EMIT, ABSORB, OR REFLECT LIGHT, WHICH IS WHY IT IS "DARK." IT ALSO APPEARS TO INTERACT VERY WEAKLY, IF AT ALL, WITH OTHER FUNDAMENTAL FORCES LIKE THE ELECTROMAGNETIC AND STRONG NUCLEAR FORCES. SOME CANDIDATES, LIKE WIMPS, ARE THEORIZED TO HAVE A WEAK INTERACTION WITH ORDINARY MATTER.

Q: IS DARK MATTER MADE OF PROTONS AND NEUTRONS?

A: NO, DARK MATTER IS UNDERSTOOD TO BE NON-BARYONIC. THIS MEANS IT IS NOT MADE OF THE FAMILIAR PROTONS AND NEUTRONS THAT FORM ORDINARY MATTER. EVIDENCE FROM BIG BANG NUCLEOSYNTHESIS AND THE COSMIC MICROWAVE BACKGROUND CONSTRAINS THE AMOUNT OF BARYONIC MATTER IN THE UNIVERSE, AND IT FALLS FAR SHORT OF THE TOTAL MASS INFERRED FROM GRAVITATIONAL OBSERVATIONS.

Q: WHAT ARE THE LEADING CANDIDATES FOR DARK MATTER PARTICLES?

A: THE LEADING CANDIDATES FOR DARK MATTER PARTICLES INCLUDE WEAKLY INTERACTING MASSIVE PARTICLES (WIMPS), AXIONS, STERILE NEUTRINOS, AND PRIMORDIAL BLACK HOLES. EACH OF THESE THEORETICAL PARTICLES POSSESSES PROPERTIES THAT COULD EXPLAIN THE OBSERVED DARK MATTER PHENOMENA.

Q: HOW ARE SCIENTISTS TRYING TO DETECT DARK MATTER DIRECTLY?

A: DIRECT DETECTION EXPERIMENTS AIM TO OBSERVE THE RARE OCCASIONS WHEN A DARK MATTER PARTICLE MIGHT COLLIDE WITH AN ATOMIC NUCLEUS IN A HIGHLY SENSITIVE DETECTOR. THESE DETECTORS ARE TYPICALLY PLACED DEEP UNDERGROUND TO SHIELD THEM FROM BACKGROUND RADIATION, AND THEY ARE DESIGNED TO REGISTER THE TINY RECOIL ENERGY PRODUCED BY SUCH AN INTERACTION.

Q: WHAT IS GRAVITATIONAL LENSING, AND HOW DOES IT RELATE TO DARK MATTER?

A: GRAVITATIONAL LENSING IS THE BENDING OF LIGHT FROM DISTANT OBJECTS BY THE GRAVITY OF INTERVENING MASS. BY OBSERVING HOW LIGHT IS DISTORTED, MAGNIFIED, OR APPEARS AS MULTIPLE IMAGES, ASTRONOMERS CAN MAP THE DISTRIBUTION OF MASS. THESE MAPS CONSISTENTLY SHOW THAT THE TOTAL MASS RESPONSIBLE FOR LENSING IS FAR GREATER THAN THE VISIBLE MATTER, INDICATING THE PRESENCE OF DARK MATTER.

Q: WHY IS THE COSMIC MICROWAVE BACKGROUND (CMB) IMPORTANT FOR UNDERSTANDING DARK MATTER?

A: THE CMB IS A SNAPSHOT OF THE EARLY UNIVERSE. THE PRECISE PATTERNS OF TEMPERATURE FLUCTUATIONS WITHIN THE CMB ARE EXTREMELY SENSITIVE TO THE COMPOSITION OF THE UNIVERSE. COSMOLOGICAL MODELS THAT INCLUDE A SIGNIFICANT AMOUNT OF NON-BARYONIC DARK MATTER ARE HIGHLY SUCCESSFUL AT EXPLAINING THESE OBSERVED CMB PATTERNS, PROVIDING STRONG SUPPORT FOR ITS EXISTENCE.

Q: WHAT IS THE DIFFERENCE BETWEEN "COLD" AND "HOT" DARK MATTER?

A: THE DISTINCTION BETWEEN "COLD" AND "HOT" DARK MATTER REFERS TO THE SPEED OF THE PARTICLES IN THE EARLY UNIVERSE. COLD DARK MATTER PARTICLES WERE SLOW-MOVING AND BEGAN TO CLUMP EARLY, FACILITATING THE FORMATION OF THE HIERARCHICAL STRUCTURES WE SEE TODAY. HOT DARK MATTER PARTICLES WERE RELATIVISTIC AND WOULD HAVE SMOOTHED OUT EARLY DENSITY FLUCTUATIONS, LEADING TO A DIFFERENT STRUCTURE FORMATION SCENARIO. CURRENT EVIDENCE FAVORS COLD OR WARM DARK MATTER.

[Dark Matter And Its Properties](#)

Dark Matter And Its Properties

Related Articles

- [d-day significance for us military operations planning](#)
- [dairy free milk alternatives us](#)
- [d-day monuments normandy](#)

[Back to Home](#)